

A CHARACTERIZATION OF QUASIPOSITIVE SEIFERT SURFACES (CONSTRUCTIONS OF QUASIPOSITIVE KNOTS AND LINKS, III)

LEE RUDOLPH [†]

(Received 8 August 1989; in revised form 7 January 1991;
annotated and arXived by the author 13 November 2004)

§0. INTRODUCTION; STATEMENT OF RESULTS

LET $T_{n,n} \subset S^3$ be a fiber surface for the torus link $O\{n,n\}$; say, to be concrete,
 $T_{n,n} = \{(z,w) \in S^3: z^n + w^n \geq 0\} \subset S^3 = \{(z,w): |z|^2 + |w|^2 = 1\}$.

CHARACTERIZATION THEOREM. *A Seifert surface S is quasipositive if and only if, for some n , S is ambient isotopic to a full subsurface of $T_{n,n}$.* $\square$

This should be contrasted with the following.

LYON'S THEOREM [0]. *Any Seifert surface is ambient isotopic to a (full) subsurface of the fiber surface of $O\{n,n\} \neq O\{n,n\}$ for some n .* $\square$

Here, a *surface* is smooth, compact, oriented, and has no component with empty boundary. A *Seifert surface* is a surface embedded in S^3 . A subsurface S of a surface T is *full* if each simple closed curve on S that bounds a disk on T already bounds a disk on S . The definition of quasipositivity is recalled in §1, after a review of braided surfaces.

The “only if” statement of the Characterization Theorem is proved in §2. Some results about graphs on braided surfaces (stated, with one eye on other applications [6], in somewhat more generality than needed here) are obtained in §3, and used in §4 to prove the “if” statement of the Characterization Theorem. A conjectural extension to ribbon surfaces in the 4-disk is discussed in §5.

Remark. The relation “ S is a full subsurface of T ” is transitive, so the Characterization Theorem has the interesting corollary (applied in [5] and [7]) that *any full subsurface of any quasipositive Seifert surface is quasipositive*.

§1. BRAIDS, BRAIDED SURFACES, AND QUASIPOSITIVITY

Let $\sigma_1, \dots, \sigma_{n-1}$ be the usual generators of the n -string braid group B_n . For $1 \leq i < j \leq n$, let $\sigma_{i,j} := (\sigma_i \cdots \sigma_{j-2})\sigma_{j-1}(\sigma_i \cdots \sigma_{j-2})^{-1}$. A *positive* (resp., *negative*) *embedded band* in B_n is any $\sigma_{i,j}$ (resp., $\sigma_{i,j}^{-1}$). An *embedded band representation* in B_n , of length k , is a k -tuple $\mathbf{b} = (b(1), \dots, b(k))$ of embedded bands. If each $b(t)$ is positive, then $\mathbf{b}$ is

[†]Partially supported by NSF grant DMS-8801959.




Fig. 1. A braided Seifert surface $S(\mathbf{b})$, $\mathbf{b} = (\sigma_{1,2}, \sigma_{2,3}^{-1}, \sigma_{1,3})$.

quasipositive. If each $b(t)$ is some usual generator $\sigma_i = \sigma_{i,i+1}$ or inverse σ_i^{-1} , then $\mathbf{b}$ is a *braidword*. (A quasipositive braidword is customarily simply called *positive*.)

Given $\mathbf{b}$, a *braided surface* $S(\mathbf{b})$ is constructed as in Fig. 1 (cf. [2]‡). More precisely, if $b(t) = \sigma_{i(t),j(t)}^{\varepsilon(t)} \in B_n$, $1 \leq i(t) < j(t) \leq n$, $\varepsilon(t) = \pm 1$, $1 \leq t \leq k$, then $S(\mathbf{b}) \subset \mathbb{R}^3 \subset S^3$ is a Seifert surface with handle decomposition $\bigcup_{s=1}^n h_s^{(0)} \cup \bigcup_{t=1}^k h_t^{(1)}$ such that, for appropriate rectangular coordinates (x, y, z) on $\mathbb{R}^3$,

- (1) $h_s^{(0)}$ is contained in the vertical half plane $\{(s, y, z): y \geq 0, z \in \mathbb{R}\}$,
- (2) $h_t^{(1)}$ is contained in the box $\{(x, y, z): i(t) \leq x \leq j(t), y \leq 0, t-1 \leq z \leq t\}$,
- (3) $h_t^{(1)}$ joins $h_{i(t)}^{(0)}$ and $h_{j(t)}^{(0)}$ with a single half-twist of sign $\varepsilon(t)$.

According to [2], this construction has a converse: if S is any Seifert surface, then there exists $\mathbf{b}$ (highly nonunique!) such that S is ambient isotopic to $S(\mathbf{b})$. S is *quasipositive* if some such $\mathbf{b}$ is quasipositive.

Note that (after enlarging its 0-handles in their half-planes as necessary) the link $\partial S(\mathbf{b})$ is presented, with respect to a horizontal braid axis (omitted from the figure) as the closed braid $\widehat{\beta}(\mathbf{b})$, where $\beta(b) := b(1) \cdots b(k)$.

§2.

The “only if” statement in the Characterization Theorem follows immediately from 2.1 and 2.2, plus the observation that the relation “ S is a full subsurface of T ” is transitive.

LEMMA 2.1. *If $\mathbf{p}$ is a positive braidword in B_m , then, for some n , $S(\mathbf{p})$ is ambient isotopic to a full subsurface of $T_{n,n}$.*

Proof. Let ∇_n be the positive braidword of length $n(n-1)$ in B_n with $\nabla(s) = \sigma_d$ if $s = (n-1)c + d$, $0 \leq c \leq n-1$, $1 \leq d \leq n-1$. Then $\widehat{\beta}(\nabla_n)$ is a torus link $O\{n, n\}$ of type (n, n) , and $S(\nabla_n)$ is a fiber surface $T_{n,n}$ (cf. [8]). Let n be the greater of m and the length of the braidword $\mathbf{p}$; make the usual identification of B_m with a subgroup of B_n . By lavishly inserting letters into $\mathbf{p}$, pad it out to ∇_n . Now $S(\mathbf{p})$ may be constructed as a (manifestly full) subsurface of $S(\nabla_n)$. $\square$

LEMMA 2.2. *If S is a quasipositive Seifert surface, then S is ambient isotopic to a full subsurface of $S(\mathbf{p})$ for some positive braidword $\mathbf{p}$ in some B_n .*

‡ The orientation conventions in [2]–[4] are opposed to those of [5], [6], and the present paper.




Fig. 2.

Proof. We may assume that $S = S(\mathbf{b})$, where $\mathbf{b} = (\sigma_{i(1), j(1)}, \dots, \sigma_{i(k), j(k)})$ is a quasipositive embedded band representation in some B_n . By inspection (cf. the 1-handle pictured in Fig. 2, $S(\mathbf{b})$ is ambient isotopic to a (full) subsurface of $S(\mathbf{p})$ where $\mathbf{p}$ is the positive braidword $(\sigma_{i(1)}, \sigma_{i(1)+1}, \dots, \sigma_{j(1)-1}, \sigma_{i(2)}, \dots, \sigma_{j(2)-1}, \dots, \sigma_{i(k)}, \dots, \sigma_{j(k)-1})$. $\square$

§3. GRAPHS ON BRAIDED SURFACES

A *graph* is a polyhedron G of dimension ≤ 1 . If $g \in G$, then g is an *isolated point* (resp., *endpoint*; *ordinary point*; *intrinsic vertex*) if the link of g in G consists of 0 (resp., 1; 2; $v(g) \geq 3$) points. G is *trivalent* if $v(g) = 3$ for each intrinsic vertex $g \in G$.

A graph E contained in a vertical half-plane $\{(s, y, z): y \geq 0, z \in \mathbb{R}\}$ is a *comb* if E is the union of a vertical interval $\{(s, y, z): y = y_0, z_1 \leq z \leq z_m\}$ and $m \geq 2$ horizontal intervals $\{(s, y, z): x = s, y_0 \geq y \geq 0, z = z_h\}, y_0 > 0, z_1 < \dots < z_m$; E is trivalent, with m endpoints (on the boundary of the half-plane) and $m - 2$ intrinsic vertices.

Let S be a surface in which a graph G is embedded. Let $N_S(G)$ be a regular neighborhood of G . G is *full* if no simple closed curve $C \subset G$ bounds a disk on S (so G is full if and only if $N_S(G)$ is), and G is *proper* if $G \cap \partial S$ is precisely the set of endpoints of G . If $N \subset S$ is a subsurface such that each component of $N \cap \partial S$ is an arc, then N is a regular neighborhood $N_S(G)$ of some proper trivalent graph $G \subset S$.

Let G' be another graph embedded in S . G and G' are *isotopic* if there exists a piecewise-smooth isotopy of S carrying G onto G' . (It may not be possible to find such an isotopy which is smooth near intrinsic vertices.) G and G' are *Whitehead-equivalent* if $N_S(G)$ and $N_S(G')$ are ambient isotopic on S .

Given a handle decomposition $S = \bigcup_{s=1}^n h_s^{(0)} \cup \bigcup_{t=1}^k h_t^{(1)}$, set $G^{(0)} := G \cap \bigcup_{s=1}^n h_s^{(0)}$, $G^{(1)} := G \cap \bigcup_{t=1}^k h_t^{(1)}$. G is *well-placed* if each point of $G^{(1)}$ is an ordinary point of G , and each component of $G^{(1)}$ is isotopic to a core arc of some $h_t^{(1)}$. If $S = S(\mathbf{b})$ is braided and $\bigcup_{s=1}^n h_s^{(0)} \cup \bigcup_{t=1}^k h_t^{(1)}$ is its given handle decomposition, G is *combed* if it is well-placed and $G^{(0)}$ is a disjoint union of combs and isolated points.

LEMMA 3.1. *Any graph on S is isotopic to a well-placed graph.*

Proof. Obvious. $\square$

LEMMA 3.2. *Any trivalent, full, proper graph on $S(\mathbf{b})$ is isotopic, by an isotopy supported on the 0-handles, to a combed graph.*

Proof. Let G satisfy the hypotheses. By 3.1, we can assume G is well-placed. Let $S(\mathbf{b}) \cap \{(s, y, z): y \geq 0, z \in \mathbb{R}\} = h_s^{(0)}$ be a 0-handle. By fullness of G and the Jordan Curve Theorem, each component of $G \cap h_s^{(0)}$ is an isolated point or a tree. By the properness of G , if e is an endpoint of $G \cap h_s^{(0)}$, then $e \in \partial h_s^{(0)}$, and e is interior to an attaching arc of some 1-handle (resp., exterior to all the attaching arcs) iff e is an ordinary point (resp., an endpoint) of G . Thus after a preliminary isotopy supported in a collar of $\partial h_s^{(0)}$, we can assume that every endpoint of $G \cap h_s^{(0)}$ is an interior point of the interval $J := S(\mathbf{b}) \cap \{(s, 0, z): z \in \mathbb{R}\}$. For each tree component E of $G \cap h_s^{(0)}$, let $I \subset J$ be the smallest subinterval containing all the endpoints of E , let A be the subarc of E joining the endpoints of I , and let $D(E)$ be the subdisk of $h_s^{(0)}$ bounded by $I \cup A$. The set of disks $D(E)$ is partially ordered by inclusion. Using trivalence of G , we can construct the desired isotopy by induction over this poset. $\square$

LEMMA 3.3. *Let $G \subset S(\mathbf{b})$ be combed. If for some s, t , there is a comb $E \subset G \cap h_s^{(0)}$ which has two or more endpoints in the attaching arc $h_s^{(0)} \cap h_t^{(1)}$, then there is a combed graph G' Whitehead-equivalent to G such that $G'^{(1)}$ has fewer components than $G^{(1)}$.*

Proof. Let $E = \{(s, y_0, z): z_1 \leq z \leq z_m\} \cup \bigcup_{i=1}^m \{(s, y, z): y_0 \geq y \geq 0, z = z_i\}$ be such a comb. Then, for some i , adjacent endpoints $(s, 0, z_i), (s, 0, z_{i+1})$ of E are in $h_t^{(1)}$. Altering G in a neighborhood of $h_t^{(1)} \cup \{(s, y, z): 0 \leq y \leq y_0, z_i \leq z \leq z_{i+1}\}$, we can replace G with a Whitehead-equivalent graph G'' which is still trivalent, full, proper, and well-placed, such that $G''^{(1)}$ has fewer components than $G^{(1)}$. (Cf. Fig. 3; the “Whitehead move” we use is the equivalent, on the level of graphs, of a handle-slide on the level of regular neighborhoods.) Then we apply 3.2 to G'' to obtain G' . $\square$




Fig. 3.

§4.

At this point, $S(\nabla_n)$ is no longer the best choice of a braided surface ambient isotopic to $T_{n,n}$. Instead, for $v := (n-1)^2 + 1$, let $\mathbf{q}$ be the quasipositive band representation of length $2(v-1)$ in B_v defined as follows: for $1 \leq s \leq v-1$, set $q(s) = \sigma_{1, v-s+1}$; for $v \leq s \leq 2(v-1)$, if $s-v = (n-1)c + d$, $0 \leq c, d \leq n-2$, then set $q(s) = \sigma_{1, v-c-(n-1)d}$. (The case $n = 3$ is illustrated in Fig. 4.)




Fig. 4.

LEMMA 4.1. $S(\mathbf{q})$ is ambient isotopic to $T_{n,n}$.

Proof. First show that $\hat{\beta}(\mathbf{q})$ is ambient isotopic to $O\{n, n\}$ (this is a straightforward exercise in braid relations and Markov moves, given that $O\{n, n\} = \hat{\beta}(\nabla_n)$). Then calculate

$$\chi(S(\mathbf{q})) = v - 2(v - 1) = 2 - v = 1 - (n - 1)^2 = n - n(n - 1) = \chi(T_{n,n}).$$

Finally, use the familiar fact that up to ambient isotopy a fiber surface is the unique Seifert surface of maximal Euler characteristic for its boundary. $\square$

In addition to its given handle decomposition $S(\mathbf{q}) = \bigcup_{s=1}^v h_s^{(0)} \cup \bigcup_{t=1}^{2(v-1)} h_t^{(1)}$, which will now be called *fine*, we need a *coarse* handle decomposition with a single 0-handle $H_1^{(0)} := h_1^{(0)}$ and $v - 1$ 1-handles $H_s^{(1)} := h_s^{(1)} \cup h_{v-s+1}^{(0)} \cup h_{v+s'}^{(1)}$ where $1 \leq s = 1 + c + (n - 1)d \leq v - 1$, $0 \leq c, d \leq n - 1$, $s' = d + (n - 1)c$. Note that, if G is well-placed with respect to the coarse handle decomposition, then G is certainly well-placed with respect to the fine one.

THEOREM 4.2. Any full subsurface of $T_{n,n}$ is quasipositive.

Proof. Let $S \subset S(\mathbf{q})$ be a full subsurface. We may assume $S \cap \partial S(\mathbf{q}) = \emptyset$. Among all graphs $G \subset S(\mathbf{q})$ with $N_{S(\mathbf{q})}(G)$ isotopic to S on $S(\mathbf{q})$, such that G is proper, full, trivalent, and well-placed with respect to the coarse handle decomposition, let G_0 be such that $G_0^{(1)}$ (also with respect to the coarse handle decomposition) has the minimal number of components. Now apply 3.2, with respect to the fine handle decomposition, to find a combed graph G_1 isotopic to G_0 . Let $E \subset G_1$ be a comb; then E has no more than one endpoint in any attaching arc of a 1-handle of the fine handle decomposition, for otherwise 3.3 would contradict the assumed minimality of G_0 .

We are nearly done. Of course $N_{S(\mathbf{q})}(G_1)$ is isotopic to S on $S(\mathbf{q})$. On the other hand, Fig. 5 shows how to perform local moves (in the vicinity of each coarse 1-handle) which effect an ambient isotopy of $N_{S(\mathbf{q})}(G_1)$ in $\mathbb{R}^3$ that pushes $N_{S(\mathbf{q})}(G_1)$ off $S(\mathbf{q})$ and onto a quasipositive braided surface. (The fact that the coarse decomposition has a single




Fig. 5. The black surface on the left is $N_{S(q)}(G_1)$; it is isotopic, in $\mathbb{R}^3$, to the black surface on the right, which can be braided by expanding the thin vertical pieces into fat 0-handles.

0-handle ensures that the local moves do not interfere with each other. It was for this reason that the coarse decomposition was introduced.) $\square$

Taken with §2, 4.2 completes the proof of the Characterization Theorem.

§5. A CONJECTURAL GENERALIZATION

Let $d : \mathbb{C}^2 \rightarrow [0, \infty[: (z, w) \mapsto (|z|^2 + |w|^2)^{1/2}$. A smoothly embedded surface $S \subset D^4(r) := d^{-1}([0, r])$, $r > 0$, is *ribbon-embedded* if $\partial S = S \cap \partial D^4(r)$ and the restriction $d^2|_S$ is a Morse function which has no local maxima on $\text{Int } S$. If $S \subset D^4(r)$ is ribbon-embedded and $r' \in]0, r[$ is a regular value of $d^2|_S$, then $S' := S \cap D^4(r')$ is ribbon-embedded in $D^4(r')$, and S' is full on S ; call (S, S') a *ribbon-embedded pair*. A surface ambient isotopic to a ribbon embedded surface is a *ribbon*; a subsurface $S' \subset S$ of a ribbon is a *subribbon* if the pair (S, S') is ambient isotopic to a ribbon-embedded pair.

Let $\Gamma_n(\epsilon)$ be the complex plane curve $\{(z, w) \in \mathbb{C}^2 : z^n + w^n = \epsilon\}$. Then, for all sufficiently small $\epsilon \neq 0$, the sets $\Gamma_n(\epsilon) \cap D^4(1)$ are mutually isotopic ribbon-embedded surfaces. Let $\tilde{T}_{n,n}$ denote any one of them.

CONJECTURE. *A ribbon is quasipositive if and only if, for some n , it is ambient isotopic to a subribbon of $\tilde{T}_{n,n}$.*

Here, by appealing to results in [2], we can define *quasipositive ribbon* as follows (the original definition involves braids, and “band representations” more general than the embedded band representations used in this paper). Let D denote the bidisk $\{(z, w) : |z| \leq 1, |w| \leq 1\}$, $\partial_1 D := \{(z, w) : |z| = 1, |w| \leq 1\}$. Let Γ be a non-singular complex plane curve which intersects $\partial_1 D$ transversely and the rest of ∂D not at all. Let $\eta : D \rightarrow D^4(1)$ be a smoothing (a homeomorphism which is a diffeomorphism except along the corner torus $\partial(\partial_1 D)$). Then $\eta(\Gamma \cap D)$ is a quasipositive ribbon, and every quasipositive ribbon arises this way.

The following facts are known, cf. [1], [2], [3]. (1) One may “push in” the interior of a Seifert surface in $S^3 = \partial D^4(1)$ to obtain a ribbon with the same boundary; if the Seifert surface is quasipositive, so is the ribbon. (2) In particular, $\tilde{T}_{n,n}$ itself is a quasipositive ribbon, for it is the push-in of $T_{n,n}$ (in fact, they are the two versions—one in the 4-disk, the other in the 3-sphere—of the Milnor fiber of $z^n + w^n$). (3) If S is a quasipositive ribbon, then for some n , S is a subribbon of $\tilde{T}_{n,n}$. (This can be seen as follows. Realize S by $\eta(\Gamma \cap D)$ as above. Without loss of generality, the completion of Γ in $\mathbb{CP}^2 \supset \mathbb{C}^2$ is non-singular and transverse to the line at infinity. Let $f(z, w)$ be the defining polynomial of Γ . For all sufficiently large $R > 0$, $\eta(\Gamma \cap D)$ is isotopic to a union of components of the ribbon-embedded surface $\{(z, w) \in \mathbb{C}^2: f(z, Rw) = 0\} \cap D^4(1)$ in $D^4(1)$. This is in turn a subribbon of $\{(z, w) \in \mathbb{C}^2: f(z, Rw) = 0\} \cap D^4(r)$ in $D^4(r)$ for generic $r \geq 1$. Because Γ is transverse to the line at infinity, $\{(z, w) \in \mathbb{C}^2: f(z, Rw) = 0\} \cap D^4(r)$ is isotopic to $\{(z, w) \in \mathbb{C}^2: z^n + w^n = 1\} \cap D^4(r)$ for all sufficiently large r , where n is the degree of f . Finally, by rescaling, this last ribbon is isotopic to $\tilde{T}_{n,n}$.)

Thus what remains conjectural is that every subribbon of $\tilde{T}_{n,n}$ is quasipositive. If this is true, then (up to ambient isotopy) an oriented link L in $S^3 = \partial D^4(r)$ bounds a piece of complex plane curve in $D^4(r)$ if and only if L has some representation as the closure of a quasipositive braid. The Characterization Theorem gives some hope that this is so, but it seems likely that entirely different techniques will be needed for a proof. ①

References

0. H. LYON: Torus knots in the complements of links and surfaces, *Mich. Math. J.* **27** (1980), 39–46.
1. L. RUDOLPH: Algebraic functions and closed braids, *Topology* **22** (1983), 191–202.
2. L. RUDOLPH: Braided surfaces and Seifert ribbons for closed braids, *Comment. Math. Helv.* **58** (1983), 1–37.
3. L. RUDOLPH: Constructions of quasipositive knots and links, I, *Nœuds, Tresses, et Singularités* (ed. C. Weber), *L'Ens. Math.* (1983), 233–245.
4. L. RUDOLPH: Constructions of quasipositive knots and links, II, *Contemp. Math.* **35** (1984), 485–491.
5. L. RUDOLPH: Quasipositive annuli (Constructions of quasipositive knots and links, IV), preprint (1990). ②
6. L. RUDOLPH: Quasipositive plumbing (Constructions of quasipositive knots and links, V), preprint (1990). ③
7. L. RUDOLPH: Quasipositivity and new link invariants, *Revista Matemática de la Universidad Complutense de Madrid* **2** (1989), 85–109.
8. J. R. STALLINGS: Constructions of fibred knots and links, *Proc. Sympos. Pure Math.* XXXII, Part 2 (Providence: AMS, 1979), 55–60.

Department of Mathematics and Computer Science
 Clark University
 Worcester
 Massachusetts 01610
 U.S.A.

ADDENDA

Typographical errors in the original publication have been corrected without notice; it is to be hoped that no new ones have been introduced. The following notes provide updates on various points.

(1) Boileau and Orevkov [10] have proved that, indeed, up to ambient isotopy an oriented link L in S^3 bounds a piece of complex plane curve in D^4 if and only if L has some representation as the closure of a quasipositive braid. In fact their proof shows that many subribbons of $\tilde{T}_{n,n}$ are quasipositive; as far as I am aware, the Conjecture in full generality remains open (but is perhaps of somewhat less interest).

(2) [5] was published as [11].

(3) A considerably expanded version of [6] was published as [12].

Additional References

10. MICHEL BOILEAU and STEPAN YU. OREVKOV: Quasipositivité d'une courbe analytique dans une boule pseudo-convexe, *C. R. Acad. Sci. Paris* **332** (2001), 825–830. MR1836094
11. L. RUDOLPH: Quasipositive annuli (Constructions of quasipositive knots and links, IV), *J. Knot Theory Ramifications* **1** (1992), 451–466. MR1194997
12. L. RUDOLPH: Quasipositive plumbing (Constructions of quasipositive knots and links, V), *Proc. Amer. Math. Soc.* **126**, 257–267. MR1452826